

GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

TDDC Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine
and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G.
Edwards F. Coles, M.D., F.A.C.P., F.A.C.G.
Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G.
Carlos A Neghme, M.D., F.A.C.G.

Shannon Marek, M.D.
Steven Pearce, M.D.
John Cluley, M.D.
Lori Hale, PA-C, MPAS

COLONOSCOPY MoviPrep

PLEASE READ ALL INSTRUCTIONS CAREFULLY!


If you need to cancel your appointment, please contact the office at least Three (3) days prior to your procedure. If the office does not receive cancellation 24 hours prior to your procedure a *cancellation fee will be charged*.

*****If the prep is not completed appropriately, your procedure may be cancelled*****

You have been scheduled for a colonoscopy on _____ at _____ .
Please arrive at facility by _____ .

- Bring your **Insurance Card** and **payment** for necessary co-pays
- Wear loose-fitting comfortable clothing. NO lipstick, contact lenses, makeup.
- **Leave Jewelry at home**
- Bring phone and fax numbers of referring physicians you would like reports sent to
- **YOU MUST HAVE A RESPONSIBLE ADULT TO DRIVE YOU HOME**
 - **Your driver must stay during your procedure.** Plan for someone to be with you 4-6 hours after your procedure as well.
 - Your procedure **WILL BE CANCELLED** if you do not have a driver to take you home.

MEDICATION GUIDELINES



- You may take your other necessary medications as you usually would
- On day of the exam, NO diuretics should be taken (Lasix, Aldactone, HCTZ, etc)
- Diabetic medications require special instructions
- **BRING A LIST OF YOUR CURRENT MEDICATIONS** (including over the counter meds)
- **If you are a female, you must notify someone if there is any possibility you maybe pregnant.**

5 days prior to your procedure (_____)

- STOP TAKING: Iron pills, Coumadin, Plavix, Aggrenox, Pletal, Effient, Pradexa, Xarelto, Eliquis etc. (unless otherwise instructed)

GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

TDDC Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine
and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G.
Edwards F. Coles, M.D., F.A.C.P., F.A.C.G.
Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G.
Carlos A Neghme, M.D., F.A.C.G.

Shannon Marek, M.D.
Steven Pearce, M.D.
John Cluley, M.D.
Lori Hale, PA-C, MPAS

-
- Avoid eating any raw fruits, vegetables or nuts(COOKED OR CANNED VEGETABLES AND FRUITS ARE ALLOWED)

IF YOU HAVE NOT HAD A BOWEL MOVEMENT 1-2 DAYS PRIOR TO YOUR EXAM, TAKE 1 BOTTLE OF MAGNESIUM CITRATE (which is sold over the counter) AT 8PM ON

1 day prior to your procedure ()

- **FOR BREAKFAST:** You may have a full liquid diet (milk, plain oatmeal, plain yogurt, pudding, nut free ice cream, cream soups)
- **REST OF THE DAY STARTING AT 12 NOON: CLEAR LIQUID DIET ONLY. NO SOLID FOODS**
 - Strained Fruit Juices without pulp (apple, White Grape, Orange or Lemonade) No red/purple juices.
 - Water (drink at least 64 ounces through the course of the day)
 - Clear broth or bouillon, coffee or tea, (without milk or non-dairy creamer)
 - All of the following as long as they are **NOT** colored **RED** or **PURPLE**
 - Gatorade- Snapple
 - Carbonated and non-carbonated soft drinks (7-up, sprite, ginger ale)
 - Kool-Aid (or other fruit flavored drinks)
 - Plain Jell-O (preferably clear or yellow, without added fruits or toppings)
- **AT 5:30PM:** Take the first dose of Reglan (generic name: Metoclopramide) to prevent nausea
- **AT 6:00PM:** Take first dose of **MOVIPREP** laxative
 - Empty 1 Pouch "A" and 1 pouch "B" into the disposable container.
 - Add lukewarm drinking water to the top line of the container. Mix to dissolve. If preferred, mix solution ahead of time and refrigerate prior to drinking. The reconstituted solution should be used within 24 hours.
 - (OPTIONAL) You can mix Crystal Light (no red flavors) with the solution for flavor
 - The container is divided into 4 marks. Every 15 minutes, drink solution down to the next mark (approximately 8 oz), until full liter is consumed. **You must then drink 32 oz of water after completing solution.**
 - (OPTIONAL) Flushable moist wipes may prevent anal irritation from excessive wiping with the prep

Day of your procedure

- **6.5 HOURS PRIOR** (at): Take the 2nd dose of Reglan (generic name: Metoclopramide) to prevent nausea
- **6 HOURS PRIOR** (at): Take second dose of **MOVIPREP** laxative

GASTROENTEROLOGY OF SAN MARCOS/NEW BRAUNFELS

TDDC Texas Digestive Disease Consultants

Diplomates of the American Board of Internal Medicine
and Subspecialty Board of Gastroenterology

W. Kern Deschner, M.D., F.A.C.G.
Edwards F. Coles, M.D., F.A.C.P., F.A.C.G.
Thomas Kepczyk, M.D., F.A.C.P., F.A.C.G.
Carlos A Neghme, M.D., F.A.C.G.

Shannon Marek, M.D.
Steven Pearce, M.D.
John Cluley, M.D.
Lori Hale, PA-C, MPAS

- Empty 1 Pouch "A" and 1 pouch "B" into the disposable container and mix as per previous instructions. Again, can also refrigerate and mix with Crystal Light for flavor.
- The container is divided into 4 marks. Every 15 minutes, drink solution down to the next mark (approximately 8 oz), until full liter is consumed. **You must drink 16 oz of water after completing solution.**
-
- **4 HOURS PRIOR (at): NO FURTHER LIQUIDS BY MOUTH(no gum,candy,ice or water) (except a sip of water with blood pressure medication only) Hold all other medications until after your procedure.**

***** If you eat or drink within 4 hours of the procedure, we may have to cancel the procedure.*****

If you have any questions please call: Barbara ext 2205 or Julie ext 2228

YOU ARE SCHEDULED AT THE FOLLOWING:

- Central Texas Medical Center** (call 512-753-3665) 1-2 days prior to exam
1301 Wonder World Drive, San Marcos, TX. 78666
- Christus Santa Rosa Out Patient Surgery-San Marcos**
Enter through "Main Entrance" of Hospital (Central Texas Medical Center) (call 830-643-8600) 1-2 prior to exam
1301 Wonder World Drive Suite 105, San Marcos, TX. 78666
- Christus Santa Rosa Out Patient Surgery-New Braunfels** (call 830-643-8600) 1-2 days prior to exam
1738 E. Common Street, New Braunfels, TX 78130
(YOU ARE SCHEDULED AT SURGERY CENTER NOT HOSPITAL)
- Seton Edgar B. Davis Admission Office** (call 830-875-7000) 1-2 days prior to exam
130 Hays Street, Luling, TX. 78648
- Resolute Hospital** (830) 500-6000 **(No pre-registration required)**
555 Creekside Way, New Braunfels, TX 78130

- For questions regarding anesthesia call **1-866-647-3631**
- For questions regarding physicians bill call **214-623-6230**